

22 PREVALENCE OF VITAMIN D DEFICIENCY IN FEMALES OF HIGHER SOCIOECONOMIC CLASS OF AHMEDABAD, GUJARAT, INDIA Authors Jayesh Vaza¹, Piyush Damor¹, Samip Sheth¹

¹Department of Orthopedics, AMC MET Medical College, Maninagar Ahmedabad, Gujarat, India.

Correspondence to: Piyush Damor, E-mail: piyusd91@gmail.com

ABSTRACT

Background: Vitamin D is a group of fat-soluble secosteroids responsible for intestinal absorption of calcium &

phosphate-. Vitamin D is a major player in calcium homeostasis and bone metabolism. In humans, the most -important compounds of vitamin D are vitamin D₂ (ergocalciferol) and vitamin D₃ (cholecalciferol). Most of the vitamin D₃ is -synthesized in skin when it is exposed to ultraviolet B rays from sunlight. Vitamin D deficiency causes osteomalacia in adults and rickets in children. Deficiency of vitamin D results in impaired bone mineralization and bone damage, which leads to bone-softening diseases.

Objective: To find out the prevalence of vitamin D deficiency in females of higher socioeconomic population of Ahmedabad, Gujarat, India.

Materials and Methods: This is a cross-sectional study, carried out among the higher socioeconomic female population in Ahmedabad City, Gujarat, India. In this study, all the 300 subjects included were tested for serum 25(OH)D level by the electro-chemiluminescence immunoassay (ECLIA) method using an automated clinical chemistry analyzer (Elecsys 2010; Roche Diagnostics)

Results: In this study, majority (45%) of the subjects were severely deficient, while 39.67% were moderately deficient.

Hence, more than 80% of the subjects were moderate to severely deficient.

Conclusion: Serum vitamin D level estimation must be done in health-care management of skeletal disorders and nutritional- review status. Serum 25(OH)D is the best measure of vitamin D exposure indicating both the effects of diet and sunlight. Severity of vitamin D deficiency also increases with the age.

KEY WORDS: Cholecalciferol, vitamin D₃, 25(OH)D

INTRODUCTION:

Vitamins are organic compounds that are required in minute quantities to sustain life. Vitamin D comprises a group of fat-soluble secosteroids responsible for intestinal absorption of calcium and phosphate. Vitamin D was

long known to be a major player in calcium homeostasis and bone metabolism. Vitamin D was first discovered at the beginning of the 20th century as a missing nutrient in children with severe bone demineralization, a disease called rickets.

In humans, the most important related compounds of vitamin D are vitamin D₂ (ergocalciferol) and vitamin D₃ (cholecalciferol).^{1,2} Vitamin D₂ is synthesized by plants (mainly mushrooms and yeast), whereas vitamin D₃ is synthesized in skin when it is exposed to ultraviolet B rays from sunlight or from artificial food source such as fatty fish, eggs, beef liver, and mushrooms. So, it is also called "sunshine vitamin."²

Vitamin D deficiency is common in females especially Muslim population in pregnancy. Vitamin D has been postulated to have a role not only calcium/phosphate homeostasis but in the prevention of cancer, autoimmune conditions, and

cardiovascular disease.³ Deficiency of vitamin D results in impaired bone mineralization and bone damage, which leads to bone-softening diseases.⁴ Individuals obtain vitamin D either through consuming vitamin D-rich foods (oily fish and dairy products) or through the skin's exposure to ultraviolet B radiation from sunlight.

This vitamin D, once it enters the body, is extracted by the liver and converted to calcidiol, which is also known as 25-hydroxy vitamin D [25(OH)D]. This 25(OH)D circulates in the blood stream and is the specific vitamin D

metabolite, measurement of which in serum helps in determining a person's vitamin D status.⁵ Kidneys convert some amount of calcidiol to calcitriol, which is the biologically active form of vitamin D and known as 1,25-dihydroxy vitamin D [1,25(OH)₂D]. Calcitriol, circulating in the blood as a hormone, regulates the concentrations of calcium and phosphate in the bloodstream and promotes the healthy growth and remodeling of bone. The best indicator of total body vitamin D storage is 25(OH)D because its half-life (2–3 weeks) is far greater than that of 1,25(OH)₂D (8–12 h).⁶ The circulating concentration of 25(OH)D is in nanogram per milliliter, almost 1000-fold higher than the concentrations of 1,25(OH)₂D. The 25(OH)D also stimulates the vitamin D receptors.

Apart from its bone health-specific use, vitamin D is also used in patients with cardiovascular disease,^{7,8} multiple sclerosis,⁹ pregnancy to prevent gestational diabetes, preeclampsia, and small infants,¹⁰ tuberculosis,¹¹ HIV,¹² and deadly form of breast cancer.¹³ Reduced levels of vitamin D in blood are related to increased mortality,¹⁴ and providing vitamin D₃ as supplementary to elderly women in institutional care has been found to decrease the risk of death. An excess of vitamin D causes abnormally high blood concentrations of calcium (hypercalcemia), which can cause overcalcification of the bones, soft tissues, heart, and kidneys. It can also damage the kidney and produce kidney stones. In addition, hypertension can result.¹⁵ The symptoms of hypervitaminosis D appear several months after administration of overdoses of vitamin D, which are dehydration, vomiting, decreased appetite, irritability, constipation, fatigue, and muscle weakness. If the sunlight exposure is not adequate, then the commonly recommended daily intake of vitamin D will not be suffice.¹⁶ According to the United States Institute of Medicine,¹⁷ the recommended daily dietary allowances of vitamin D are: up to 1 year, 400 IU; from 1 year to 70 years, 600 IU; and after 70 years, 800 IU (conversion: 1 µg = 40 IU and 0.025 µg = 1 IU)

MATERIALS AND METHOD

This is a cross-sectional study, carried out among the higher socioeconomic populations of females in Ahmedabad City, Gujarat, India, during April 1, 2015, to September 30, 2016. In this study, all the 300 subjects who came for the estimation of 25(OH)D level have been included. Informed consent was obtained from all the subjects. The exclusion criteria include: (i) pregnant and lactating women and (ii) those who were taking vitamin supplements. Blood samples were collected from all the 300 subjects after overnight fasting and under aseptic pre-caution into plain vacutainers and labeled properly. After that, serum was separated and labeled properly. Then, the serum 25(OH)D level estimation was done by the electro-chemiluminescence immunoassay (ECLIA) method using an automated clinical chemistry analyzer (Elecsys 2010; Roche Diagnostics).

Reference range for the serum 25(OH)D level recommended by Mayo Medical Laboratory is 29–80 ng/mL.¹⁸ Mayo Medical Laboratories¹⁸ has categorized serum vitamin D level < 10 ng/mL as severe deficiency, between 10 and 28 ng/mL as mild to moderate deficiency, between 29 and 80 ng/mL as optimal levels, and >80 ng/mL as toxicity possible. But, in this study, we considered the serum 25(OH)D level greater than 30 ng/dL as adequate, between 21 and 28 ng/dL as mildly deficient, between 10 and 20 ng/dL as moderately deficient, and between 0 and 9 ng/dL as severely deficient.

RESULTS

This cross-sectional study was carried out for period of 18 months, and a total of 300 patients were selected for serum 25(OH)D level estimation. Among these 300 female patients, 21 (7%) were having adequate level, 28 (8.3%) mild deficiency, 119 (39.7%) moderate deficiency, and 135 (45%) severe deficiency of vitamin D. From total 300 subjects, 28 (9.3%) were between 0 and 20 years, 76 (25.3%) 21 and 40 years, 107 (35.67%) 41 and 60 years, and 89 (29.6%) older than 60 years.

For severely deficient cases, of the total 135 cases, 4 (2.96%) subjects were aged between 0 and 20 years, 33 (24.44%) 21 and 40 years, 49 (36.3%) 41 and 60 years, and 49 (36.3%) older than 60 years.

DISCUSSION

In this study 7% were having adequate level of vitamin D, 8.3% mild deficiency, 39.6% moderate deficiency, and 45% severe deficiency of vitamin D. In general, majority of the deficient subjects (35.6%) were aged between 41 and 60 years, which is followed by older than 60 years (29.6%). So, two-third of the subjects was aged older than 40 years. The majority of the deficient subjects (45%) were severely deficient, which is followed by moderately deficient subjects (39.6%). Hence, more than 85% of the subjects were moderate to severely deficient. Severity of deficiency also increases with the age.

In the study by Arya et al.¹⁹ in 2004, of the total 92 subjects, 78.3% were having vitamin D deficiency. In the

study by Harinarayan et al.²⁰ in 2004, of the total 316 subjects, 69.3% were having vitamin D deficiency. In the study by Vupputuri et al.²¹ in 2006, of the total 105 subjects, 94.3% were having vitamin D deficiency. In the study by Zargar et al.²² in 2007, of the total 92 subjects, 83% were having vitamin D deficiency. In the study by Agarwal et al.²³ in 2013, of the total 200 subjects, 58% were having vitamin D deficiency. In this study, of the total 444 subjects, 93.3% were having vitamin D deficiency, which is closely similar to the study by Vupputuri et al. and higher than the studies by Zargar et al. and Arya et al. In one of the study of 18 cities spread all over India done by Beloyartseva et al.²⁴ of the total 2,119 subjects, 79% were found deficient in vitamin D level.

Conclusion

Vitamin D plays an important role in a wide range of physiologic functions. Serum vitamin D level estimation must be done in health-care management of skeletal disorders and nutritional review status of the population. Serum 25(OH)D is the best measure of vitamin D exposure indicating both the effects of diet and sunlight. Persons with sun-avoidance behaviors were at higher risk for developing vitamin D deficiency. Vegetarian population has higher risk of vitamin D deficiency. Severity of vitamin D deficiency also increases with the age

Acknowledgments

We authors are very much thankful to all respected patients and all technical staff members for their valuable work in our study.

References

1. Holick MF. High prevalence of vitamin D inadequacy and implications for health. *Mayo Clin Proc* 2006;81(3):353–73.
2. Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr* 2004;80(6 Suppl):1678S–88S.
3. Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357(3):266–81.
4. Grant WB, Holick MF. Benefits and requirements of vitamin D for optimal health: a review. *Altern Med Rev* 2005;10(2):94–111.
5. Hollis BW. Assessment of vitamin D nutritional and hormonal status: what to measure and how to do it. *Calcif Tissue Int* 1996;58(1):4–5.
6. Haddad JG Jr, Rojanasathit S. Acute administration of 25-hydroxycholecalciferol in man. *J Clin Endocrinol Metab* 1976;42(2):284–90.
7. Pittas AG, Chung M, Trikalinos T, Mitri J, Brendel M, Patel K, et al. vitamin d and cardiometabolic outcomes: a systematic review. *Ann Intern Med* 2010;152(5):307–14.
8. McGreevy C, Williams D. New insights about vitamin D and cardiovascular disease: a narrative review. *Ann Intern Med* 2011;155(12):820–6.
9. Ascherio A, Munger, KL, Simon, KC. Vitamin D and multiple sclerosis. *Lancet Neurol* 2010;9:599–612.
10. Aghajafari F, Nagulesapillai T, Ronksley PE, Tough SC, O'Beirne M, Rabi DM. Association between maternal serum 25-hydroxyvitamin D level and pregnancy and neonatal outcomes: systematic review and meta-analysis of observational studies. *BMJ* 2013;346:f1169.
11. Luong K, Nguyen LT. Impact of vitamin D in the treatment of tuberculosis. *Am J Med Sci* 2011;341(6):493–8.
12. Spector SA, Nguyen LT. Vitamin D and HIV: letting the sun shine in. *Top Antivir Med* 2011;19(1):6–10.
13. Grotsky DA, Gonzalez-Suarez I, Novell A, Neumann MA, Yaddanapudi SC, Croke M, et al. BRCA1 loss activates cathepsin L-mediated degradation of 53BP1 in breast cancer cells. *J Cell Biol* 2013;200:187–202.
14. Zittermann A, Gummert JF, Börgermann J. Vitamin D deficiency and mortality. *Curr Opin Clin Nutr Metab Care* 2009;12(6):634–9.
15. Dionne JM, Abitbol CL, Flynn JT. Hypertension in infancy: diagnosis, management, and outcome. *Pediatr Nephrol* 2012;27(1):17–32.
16. Glerup H, Mikkelsen K, Poulsen L, Hass E, Overbeck S, Thomsen J, et al. Commonly recommended daily intake of vitamin D is not sufficient if sunlight exposure is limited. *J Intern Med* 2000;247(2):260–8.
17. Ross AC, Taylor CL, Yaktine AL, Del Valle HB. Dietary Reference Intakes for Calcium and Vitamin D. Washington, DC: National Academies Press, 2011. p. 435.
18. Categorization of Vitamin D Levels: Mayo Medical Laboratories: Vitamin D testing (article online). Available at <http://www.mayomedicallaboratoires.com/articles/vitamind/>

index.html (last accessed March 25, 2011).

19. Arya V, Bhambri R, Godbole MM, Mithal A. Vitamin D status and its relationship with bone mineral density in healthy Asian Indians. *Osteoporos Int* 2004;15(1):56–61.
20. Harinarayan CV, Ramalakshmi T, Venkataprasad U. High prevalence of low dietary calcium and low vitamin D status in healthy south Indians. *Asia Pac J Clin Nutr* 2004;13(4):359–64.
21. Vupputuri MR, Goswami R, Gupta N, Ray D, Tandon N, Kumar N. Prevalence and functional significance of 25-hydroxyvitamin D deficiency and vitamin D receptor gene polymorphisms in Asian Indians. *Am J Clin Nutr* 2006;83(6):1411–9.
22. Zargar AH, Ahmad S, Masoodi SR, Wani AI, Bashir MI, Laway BA, et al. Vitamin D status in apparently healthy adults in Kashmir Valley of Indian subcontinent. *Postgrad Med J* 2007;83(985):713–6.
23. Agrawal NK, Sharma B. Prevalence of osteoporosis in otherwise healthy Indian males aged 50 years and above. *Arch Osteoporos* 2013;8(1–2):116.
24. Beloyartseva M, Mithal A, Kaur P, Kalra S, Baruah MP, Mukhopadhyay S, et al. Widespread vitamin D deficiency among Indian health care professionals. *Arch Osteoporos* 2012;7(1–2):187–92.